



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/12, C07K 14/47, 14/715</b>		A1	(11) International Publication Number: <b>WO 98/49297</b>
			(43) International Publication Date: 5 November 1998 (05.11.98)
(21) International Application Number: PCT/US98/07439		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 14 April 1998 (14.04.98)		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 60/044,835 25 April 1997 (25.04.97) US			
(71) Applicant: AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07940 (US).			
(72) Inventors: BINGHAM, Brendan, William; 31 Tamarack Court, Newtown, PA 18940 (US). YOUNG, Kathleen, H.; 1706 Society Place, Newtown, PA 18940 (US). WOOD, Andrew, Timothy; 2051 Leedom Drive, Newtown, PA 18940 (US). BIRSAN, Camelia; 1507 Hunter's Glen Drive, Plainsboro, NJ 08536 (US).			
(74) Agents: WALSH, Andrea, C.; American Home Products Corporation, Patent Law Dept. 2B2, One Campus Drive, Parsippany, NJ 07054 (US) et al.			
(54) Title: NEURONAL MORT1 ISOFORMS			
(57) Abstract <p>A MORT1 gene initially cloned from HeLa cells and identified as a member of the receptor mediated apoptotic pathway, is expressed in the human neuronal cell line, NTERA2. Isolation of the MORT1 from this cell line revealed a transcript isoform that differed from the known MORT1 sequence by a deletion of 21 base pairs (bp 172-192 of the coding sequence). Cloning of MORT1 from adult human brain revealed two isoforms, one similarly deleted for bp 172-192, the other with a basepair substitution, A for G at position 173. Assessment of MORT1 function in a yeast two hybrid system indicates that the deleted and intact forms of MORT1 differ in their capacity to interact with other members of the apoptotic pathway.</p>			

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## Neuronal MORT1 Isoforms

### Field of the Invention

The present invention relates to human intracellular death-domain-  
5 containing apoptosis signaling proteins. In particular, the present  
invention relates to novel human neuronal MORT1 isoforms.

### Background

Apoptosis, or programmed cell death in multicellular organisms, is  
one of the fundamental means by which a cell can respond to  
10 environmental changes. One of the best studied mammalian apoptosis  
systems involves Fas (also designated APO-1 and CD95), a type I  
membrane receptor that, when crosslinked by its cognate ligand, induces  
apoptosis in a wide variety of cells (for review, see Nagata, 1994). The  
extracellular interaction of Fas ligand with the cell membrane-spanning  
15 Fas receptor activates an intracellular signal transduction cascade finally  
activating proteases in the IL-1 $\beta$ -converting enzyme (ICE) family (Henkart,  
1996). Transduction of an apoptosis signal depends on interaction  
between the intracellular "death domain" of Fas with a cytoplasmic 23-  
kDa protein, MORT1 [(Boldin, *et al.*, 1995), also termed FADD  
20 (Chinnaiyan, *et al.*, 1995)]. The events leading from the production of an  
activated Fas trimer complex to cell destruction mediated by ICE-like  
proteases are yet to be determined, but recruitment of two MORT1/FADD  
molecules into a death-inducing signaling complex with the death domain  
of Fas appears to be a necessary step (Kischkel, *et al.*, 1995). The end  
25 result of this pathway is cell death by a distinctive mechanism  
characterized by nuclear and cytoplasmic condensation and DNA  
fragmentation.

The human MORT1/FADD gene spans approximately 3.6 kb and  
contains two exons (286 and 341 bp) separated by a 2.0-kb intron.  
30 MORT1/FADD was mapped to chromosome 11q13.3 by the independent  
techniques of PCR screening of somatic cell hybrid mapping panels and

fluorescence *in situ* hybridization (Kim, *et al.*, 1996). Knowledge of the chromosomal location and gene structure of MORT1/FADD will assist efforts to determine its involvement in genetic disorders of apoptosis. Defects in apoptosis due to mutations in the Fas receptor have been  
5 described (Fisher, *et al.*, 1995; Rieux-Laucat, *et al.*, 1995) in patients with a rare autoimmune lymphoproliferative syndrome (ALPS) including nonmalignant lymphadenopathy, hepatosplenomegaly, and expanded populations of CD3<sup>+</sup>CD4<sup>+</sup>CD8<sup>-</sup> lymphocytes. However, the existence of some patients with clinical findings of ALPS, but no Fas mutation,  
10 suggests that defects in other proteins in the Fas pathway may also produce ALPS. The functional involvement of MORT1/FADD in the Fas pathway makes it a candidate for mutation analysis in ALPS.

Localization of MORT1/FADD to 11q13.3 also makes it a candidate for human diseases associated with this part of the genome.  
15 IDDM4, a predisposing locus for familial insulin-dependent diabetes mellitus (IDDM) has been mapped to this region by linkage studies (Cordell, *et al.*, 1995; Davies, *et al.*, 1994; Hashimoto, *et al.*, 1995). The pathogenesis of IDDM may involve autoimmune T lymphocyte-mediated destruction of pancreatic  $\beta$  islet cells that produce insulin (Tisch and  
20 McDevitt, 1996). MORT1/FADD's chromosomal location, coupled with its known role in lymphocyte apoptosis, makes it a candidate for mutational analysis in patients with IDDM4-linked familial diabetes.

In addition, the 11q13 region is amplified in several human malignancies, including carcinoma of the breast, bladder, esophagus, head  
25 and neck, and lung (Schuuring, 1995; Szepetowski, *et al.*, 1995). Amplification of this region has been associated with poor prognosis in patients with operable breast cancer (Schuuring, *et al.*, 1992). The mapping of MORT1/FADD to the amplicon of the breast cancer cell line MDA-MB-134-VI (Lafage, *et al.*, 1992) raises the possibility of its  
30 involvement in tumor growth. Future linkage and mutation analysis of MORT1/FADD in other diseases may support the hypothesis that

disregulation of cell death is a fundamental mechanism for the pathogenesis of human disease (Thompson, 1995).

A single gene can encode more than one mRNA transcript by transcriptional processing events such as use of alternative promoters, alternative splicing, and alternative polyadenylation (Farrow, 1997; Lewin, 1994). The resulting variant transcript isoforms may differ in stability, translatability, or protein sequence encoded, each of which may impact upon the function of the encoded protein. Transcript variants resulting from alternative RNA processing can be tissue-specific, developmentally regulated, endocrine regulated, or can appear in response to specific exogenous cues. In addition, transcript variants of a gene can result from differences in genomic sequence (among individuals or among cell lines). To date, only one transcript isoform has been reported for the MORT1/FADD gene (Boldin, et al., 1995; Chinnaiyan, et al., 1995). This invention describes the identification of two new isoforms of MORT1.

#### Summary of the Invention

The present invention is based on the discovery of novel transcript isoforms of the human neuronal MORT1 gene. The transcript isoforms and the proteins encoded by them are useful as screening agents in the diagnosis of CNS diseases and in the discovery of CNS-specific anti-apoptotic compounds. The DNA, RNA, and proteins encoded by them are useful in a "gene therapy" scenario, either as *in vivo* therapeutic agents in humans, or as experimental tools in manipulating neuronal apoptosis in cell culture and animal model systems.

Accordingly, in one embodiment, the invention is directed to a neuronal protein encoded by an MORT1 gene, wherein the gene is isolated from NTERA2 or adult human brain tissue. The cDNA encoding the neuronal protein differs from the known MORT1 gene by the deletion of 21 base pairs and by base pair substitutions.

30

### Description of the Figures

Figure 1: Investigation of the MORT1 transcript isoforms with MACH $\alpha$ 1 C360S in a yeast two-hybrid system: Activation of the HIS3 reporter gene. Each MORT1 transcript isoforms was expressed as a fusion protein with Gal4 DNA binding domain. MACH $\alpha$ 1 C360S was expressed as a fusion protein with the Gal4 activation domain. Yeast strains were plated at a cell density of  $5 \times 10^1$ ,  $5 \times 10^2$ ,  $5 \times 10^3$ ,  $5 \times 10^4$  and  $5 \times 10^5$  (left to right within a plate. Sibling isolates from yeast strains expressing a MORT1 transcript isoform with MACH $\alpha$ 1 C360S fusion proteins were plated on histidine deficient media containing 0, 20 or 40 mM 3-aminotriazole (3-AT). The top row of plates (1) contained yeast strains that express two fusion proteins as follows: YCB5: MORT1-pAS1 / MACH $\alpha$ 1 C360S -pACTII; YCB9: MORT1 $\Delta$ 21-pAS1 / MACH $\alpha$ 1 C360S -pACTII; YCB16: MORT1 G173A -pAS1 / MACH $\alpha$ 1 C360S -pACTII. The bottom row of plates (3) contains negative control strains which express only the MORT1 transcript isoform fusion protein with a recombinant vector containing nonrelated heterologous DNA, as follows: YCB5.1: MORT1 -pAS1 / SNF4 -pACTII; YCB9.1: MORT1 G173A / SNF4-pACTII; YCB16.1: MORT1 $\Delta$ 21 -pAS1 / SNF4-pACTII. Functional interaction of the MORT1 transcript isoform fusion protein with the MACH $\alpha$ 1C360S fusion protein reconstitutes the function of the Gal4 protein and drives HIS3 reporter gene activity thereby linking MORT1 isoform / MACH $\alpha$ 1C360S interaction to histidine prototrophy and yeast cell growth.

Figure 2: Investigation of the MORT1 transcript isoforms with MACH $\alpha$ 1 C360S in a yeast two-hybrid system: Activation of the CYH2 reporter gene. Each MORT1 transcript isoforms was expressed as a fusion protein with Gal4 DNA binding domain. MACH $\alpha$ 1 C360S was expressed as a fusion protein with the Gal4 activation domain. Yeast strains were plated at a cell density of  $5 \times 10^1$ ,  $5 \times 10^2$ ,  $5 \times 10^3$ ,  $5 \times 10^4$  and  $5 \times 10^5$  (left to right within a plate. Sibling isolates from yeast strains

expressing a MORT1 transcript isoform with MACH $\alpha$ 1 C360S fusion proteins were plated on selective media containing 0, 8 or 12 ug cycloheximide / mL media. The top row of plates (1) contained yeast strains that express two fusion proteins as follows: YCB5: MORT1-pAS1 / MACH $\alpha$ 1 C360S -pACTII; YCB9: MORT1 $\Delta$ 21-pAS1 / MACH $\alpha$ 1 C360S -pACTII; YCB16: MORT1 G173A -pAS1 / MACH $\alpha$ 1 C360S -pACTII. The bottom row of plates (3) contains negative control strains which express only the MORT1 transcript isoform fusion protein with a recombinant vector containing nonrelated heterologous DNA, as follows: YCB5.1: MORT1 -pAS1 / SNF4 -pACTII; YCB9.1: MORT1 G173A / SNF4-pACTII; YCB16.1: MORT1 $\Delta$ 21 -pAS1 / SNF4-pACTII. Functional interaction of the MORT1 transcript isoform fusion protein with the MACH $\alpha$ 1C360S fusion protein reconstitutes the function of the Gal4 protein and drives CYH2 reporter gene activity thereby linking MORT1 isoform / MACH $\alpha$ 1C360S interaction to sensitivity to cycloheximide and abrogation of yeast cell growth.

#### Detailed Description of the Invention

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA technology, and yeast biology and genetics, which are within the skill of the art. Such techniques are explained fully in the literature (e. g., Ausubel, *et al.*, 1993; Coico, 1994; Freshney, 1987; Glover, 1985; Griffin and Griffin, 1994; Hanes and Higgins, 1984; Perbal, 1988; Rose *et al.*, 1990; Sambrook, *et al.*, 1989).

All patents, patent applications and publications cited herein, whether supra or infra, are hereby incorporated by reference in their entirety.

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below. By "MORT1 transcript isoform" is meant a nucleic acid molecule, including DNA, RNA, mRNA, cDNA derived from the mRNA, or even synthetic

DNA, which is derived either directly or indirectly from a MORT1 genomic sequence. As used herein, the term specifically excludes the transcript described in Chinnaiyan, et al. (1995), Boldin, et al. (1995), and Genbank deposit x84709, and as Genbank x84709 MORT1. The term MORT1 $\Delta$ 21  
5 denotes a cDNA isoform isolated from NTERA2 cells (SEQ. ID NO. 1) or adult human brain (SEQ. ID NO. 3) that has a specific 21 base pair deletion (bp 172-192 of the coding sequence) as compared to the published MORT1. The term MORT1G173A (SEQ. ID NO. 5) denotes a  
10 cDNA isoform isolated from human brain that has a nucleotide substitution (A for G) at bp position 173 of the MORT1 coding sequence. Herein the A of the MORT1 translational initiation codon is designated position 1, and corresponds to bp 145 of Genbank sequence x84709.

The above terms (MORT1 transcript isoform, MORT1 $\Delta$ 21, MORT1G173A) encompass the genomic MORT1 sequence encoding  
15 them, including introns and exons. Alternatively, a transcript isoform can include transcripts of the genomic sequences which lack one or more introns or exons, or transcripts which incorporate noncoding or coding sequences different from those found in the full-length, wild-type genomic sequence. Transcript isoforms may arise by any of a number of naturally  
20 occurring processes, including but not limited to mutation, alternative initiation, alternative splicing, and alternative polyadenylation, each of which may affect the primary structure or some other aspect of the function of the encoded protein. The protein product of a transcript isoform may contain amino acids which differ from the wild-type protein  
25 due to insertion, deletion, or frameshifting of coding nucleotides. For example, the NTERA2 and the adult human brain isoforms of MORT1 are depicted in SEQ. ID NOS. 1 and 3.

The cDNA encoding intact MORT1 was generated using standard PCR techniques (Finney, 1993; Griffin and Griffin, 1994). HeLa cell cDNA  
30 prepared by standard reverse transcriptase/PCR was the source of DNA template. Additionally, MORT1 cDNA was generated using a human



placental cDNA library as the template source. In each case, a MORT1 fragment was amplified by PCR, cloned and sequenced. DNA sequences obtained from placental and HeLa tissue were compared with the MORT1 sequence in Genbank deposit x84709.

5           The existence of a human neuronal MORT1 transcript was determined by RT/PCR. Human neuronal NTERA2 cells were grown under conditions to promote terminal differentiation to the neuronal phenotype and induced to undergo apoptosis by incubation with staurosporine. RNA extracted from these cells was subjected to RT/PCR using MORT1-specific primers. The resulting MORT1 fragment was cloned and its  
10           sequence compared to the MORT1 sequence in Genbank deposit x84709.

          MORT1 was also cloned from fetal and adult human brain cDNA libraries by PCR. MORT1 fragments were amplified, cloned, and their sequence compared to the MORT1 sequence in Genbank deposit x84709.  
15           In summary, the MORT1 clones deriving from HeLa cells, the human placental library, and fetal human brain matched the Genbank MORT1 sequence. MORT1 clones from NTERA2 cells and three of five clones derived from human adult brain were deleted for bp 172-192 of the MORT1 coding sequence. The other two of the five clones derived from  
20           human adult brain had a single base pair substitution (G173A) relative to the Genbank MORT1 sequence. The 21 bp deletion and the G173A substitution both generate a Glu-Pro-Glu amino acid sequence at positions 56-58 of the protein sequence, versus the corresponding Glu-Pro-Gly sequence of the wild-type protein. In addition, the human neuronal  
25           MORT1 isoforms include some other basepair substitutions relative to Genbank sequence, as depicted in SEQ. ID NO. 1 and 3.

          Human neuronal MORT1 transcript isoforms are also cloned from human cDNA phage libraries by probing with a radiolabelled MORT1 probe using published DNA hybridization methods (Ausubel *et al.*, 1993;  
30           Sambrook, *et al.*, 1989). Many human brain sub-region libraries are commercially available (Clontech; Stratagene) for screening. For tissues

or neuronal cell lines for which no commercially available library exists, custom synthesis of a cDNA library is performed from poly(A) + RNA (service available from Clontech). Total RNA and poly(A) selection are performed according to published methods (Glover, 1985; Sambrook, *et al.*, 1989).

Human MORT1 genomic sequence is cloned using standard PCR techniques (Griffin and Griffin, 1994) from human genomic DNA prepared by standard methods (Glover, 1985; Sambrook, *et al.*, 1989).

The relative abundance of neuronal MORT1 transcript isoforms within a given tissue or cell line is assessed by RNase protection and S1 nuclease mapping, performed according to published methods (Ausubel, *et al.*, 1993; Berk and Sharp, 1977; Lee and Costlow, 1987).

Specifically, oligonucleotide probes are designed to distinguish the 21 nucleotide difference between the deleted and intact MORT1 isoforms.

A yeast two-hybrid system as described in Young and Ozenberger, WO 95/34646 published December 21, 1995, the whole of which is incorporated herein, was used to functionally characterize the ability of the MORT1 transcript isoforms to interact with MACH $\alpha$ 1 C360S, a protein component of the cytoplasmic apparatus of the Fas / APO1 and TNF receptors. Expression vectors were constructed by fusing the GAL4 DNA-binding domain to the MORT1 transcript isoforms. A second expression vector was constructed by fusing the GAL4 Activation domain to MACH $\alpha$ 1 C360S. To determine the ability of the MORT1 transcript isoforms to interact with MACH $\alpha$ 1 C360S, yeast strains were generated expressing a single MORT1 transcript isoform with MACH $\alpha$ 1 C360S, or a single recombinant plasmid encoding a fusion protein with its companion vector containing unrelated heterologous DNA. Strains were tested for productive protein-protein interaction via reporter gene activity and a change in the yeast cell phenotype.

These data suggest that the MORT $\Delta$ 21 isoform fusion protein is impaired in its ability to functionally interact with MACH $\alpha$ 1 C360S fusion

protein. The yeast strains expressing MORT $\Delta$ 21 with MACH $\alpha$ 1 C360S demonstrated decreased histidine prototrophy, and decreased cell growth, as well as decreased sensitivity to cycloheximide and increased cell growth. These were observed in comparison to the growth characteristics of yeast strain expressing the fusion proteins for MORT1 with MACH $\alpha$ 1 C360S, or MORTG173A with MACH $\alpha$ 1 C360S.

The subject nucleic acids find a wide variety of applications including use as translatable transcripts, hybridization probes, PCR primers, therapeutic nucleic acids, etc.; use in detecting the presence of genes and gene transcripts involved in apoptosis, in detecting or amplifying nucleic acids encoding additional MORT1 homologs and structural analogs, and in gene therapy applications. The gene products encoded by these transcripts have utility in serving as target proteins in the development of therapeutics for the manipulation of the apoptotic pathway.

The following experiments and examples are offered by way of illustration and not by way of limitation.

#### Examples

##### **Example 1: Cloning of MORT1 from HeLa cells and from a human placental cDNA library.**

As a control, the intact MORT1 gene was cloned as described in Boldin, *et al.*, (1995). Oligonucleotides were prepared on an ABI oligosynthesizer, designed according to the published cDNA sequence of human MORT1 (Genbank x84709). A 25 base 5' sense oligonucleotide containing a NcoI site and the start codon ("5' MORT1," 5'-ACC CCG CCA TGG ACC CGT TCC TGG T-3', corresponding to bases -8 to +17), and a 24 base 3' antisense oligonucleotide spanning the stop codon ("3' MORT1," 5'-ACG GGC CCA TCA GGA CGC TTC GGA-3', complementary to bases 636 to 613 of the MORT1 coding sequence), were synthesized. The cDNA encoding intact MORT1 was generated using standard PCR techniques (Finney, 1993). Thermal cycling of these reactions was

performed under the regime, 1 min at 95C, then 30 cycles (40 sec at 95C, 1 min at 60C, 1 min at 72C), and 10 min at 72C, using the oligonucleotides "5' MORT1" and "3' MORT1." HeLa cell cDNA and a human placental (matchmaker) cDNA library (Clontech) served as the DNA templates. In each case, a 640 bp fragment was obtained by PCR and subsequently ligated into the cloning vector pCR II (Invitrogen). The HeLa and placental MORT1 cDNA sequences matched the coding sequence of the published MORT1 sequence (Genbank deposit x84709).

**Example 2: Cloning of a neuronal MORT1 from NTERA2 cells  
by RT/PCR using a gene-specific RT primer**

In order to investigate whether MORT1 was present in neuronal cells, NTERA-2 cells, which originate from a human teratocarcinoma and represent a late embryonic human neuronal phenotype, were investigated for the presence of MORT1. NTERA2 cells (Stratagene) were grown under conditions to promote terminal differentiation to the neuronal phenotype, including long-term incubation in retinoic acid (Andrews, 1984). Eighteen hours prior to RNA extraction, staurosporine (Calbiochem) was added to the cell culture medium to a final concentration of 100 nM to induce apoptosis. Total RNA extraction was by the RNeasy Total RNA Kit (Life Technologies) using reagents and instructions provided by the manufacturer. Reverse transcription of total RNA was performed with the reagents and instructions for gene-specific reverse transcription provided in the BRL Preamplification Kit (Life Technologies). Specifically, a 27 base 3' antisense oligonucleotide ("MORT1 downstream," 5'-TAG ATG CCT GTG GTC CAC CAG CGC AAA-3', complementary to bases 663 to 637 of the MORT1 coding sequence) was synthesized on an ABI oligosynthesizer. Reverse transcription of 1 µg total RNA was primed using this oligonucleotide.

PCR amplification of NTERA2 cDNA was performed using DNA Taq polymerase and other reagents provided by the manufacturer (Life Technologies) and equimolar amounts of oligonucleotides "5' MORT1"

and "3' MORT1" (see above). Thermal cycling of the PCRs was in the Gene Amp 2400 thermal cycler (Perkin Elmer) under the regime, 1 min at 95C, 35 cycles (15 sec at 95C, 30 sec at 50C, 1 min at 72C), 7 min at 72C. The resulting PCR product (the cDNA corresponding to the

5 MORT1Δ21 isoform) was agarose gel purified using Qiaex II reagents and instructions provided by the manufacturer (Qiagen) and ligated to the pCR2.1 cloning vector (Invitrogen). Recombinant plasmids were obtained by standard methodology, including transformation into One Shot *E. coli* cells (Invitrogen), growth of transformed bacterial cells on LB-agar plates

10 containing 100 µg/mL ampicillin (Sigma), growth of bacterial colonies in LB Medium (Life Technologies,), and preparation of plasmid DNA using the QIAprep Spin Miniprep Kit (Qiagen). DNA sequence was obtained using cycle sequencing dideoxy terminator reagents (ABI) and synthetic oligonucleotides (see below) reacted in a Gene Amp 9600 thermal cycler

15 (Perkin Elmer) under manufacturer's instructions. Reactions were run and data were generated on an ABI automated sequencer. Sequence data were analyzed using Lasergene software (DNA Star).

Oligonucleotides for use in DNA sequencing were prepared (on an ABI synthesizer), designed according to the sequence of the polylinker

20 region of pCR2.1 and published MORT1 sequence. The following oligonucleotides were generated: "TA Forward" (5'-CAG GAA ACA GCT ATG ACC ATG-3', corresponding to the sense strand of the lacZ gene in pCR2.1, 67 bp upstream of the TA cloning site), "TA Reverse" (5'-ACG TTG TAA AAC GAC GGC CAG-3', corresponding to the antisense strand

25 of the lacZ gene in pCR2.1, 112 bases downstream of the TA cloning site), "MORT 150" (5'-ACC TCT TCT CCA TGC TGC TG-3', sense strand, corresponding to bases 131 to 150 of the MORT1 coding sequence), "MORT 230" (5'-TCG AAG TCG TCG ACG CGC CG-3', antisense strand, complementary to bases 248 to 229 of the MORT1 coding sequence),

30 "MORT 400" (5'-TCG ACA GCA TCG AGG ACA GA-3', sense strand, corresponding to bases 377 to 396 of the MORT1 coding sequence), and

"MORT 420" (5'-GAT TCT CAG TGA CTC CCG CA-3', antisense strand, complementary to bases 441 to 422 of the MORT1 coding sequence).

**Example 3: Cloning of MORT1 from NTERA2 cells using nested primers**

- NTERA2 cells (Stratagene) were grown under conditions to
- 5 promote terminal differentiation to the neuronal phenotype, including long-term incubation in retinoic acid (Andrews, 1984). Eighteen hours prior to RNA extraction, staurosporine (Calbiochem) was added to the cell culture medium to a final concentration of 100 nM to induce apoptosis. Total RNA extraction was by the BRL RNeasy Total RNA Kit (Life Technologies,
- 10 Inc.) using reagents and instructions provided by the manufacturer. Reverse transcription of total RNA was performed with the reagents and instructions provided in the BRL Preamplification Kit (Life Technologies, Inc.). Specifically, reverse transcription of 4 µg total RNA was primed using oligo(dT).
- 15 Oligonucleotides for use in PCR were prepared (on an ABI oligosynthesizer), designed according to the published MORT1 sequence (Genbank x84709). A 25 base 5' sense oligonucleotide ("MORT1 upstream," 5'-AAG CGG CGA GAC CTG GCC ACG GCC A-3', corresponding to bases -90 to -66, relative to the translational start site)
- 20 was synthesized. PCR amplification of NTERA2 cDNA was performed using DNA Taq polymerase and other reagents provided by the manufacturer (Life Technologies) and equimolar amounts of "MORT1 upstream" and "MORT1 downstream" (see above) oligonucleotides. Thermal cycling of the PCRs was in the Gene Amp 2400 thermal cycler
- 25 (Perkin Elmer, Inc.) under the regime, 1 min at 95C, 35 cycles (15 sec at 95C, 30 sec at 50C, 1 min at 72C), 7 min at 72C. A 2 µL aliquot of the 100 µL reaction served as the DNA template for further PCR amplification using equimolar amounts of oligonucleotides "5' MORT1" and "3' MORT1" (see above) under the same thermal cycling regime.
- 30 The resulting PCR product (the cDNA corresponding to the MORT1Δ21 isoform) was agarose gel purified using Qiaex II reagents and

instructions provided by the manufacturer (Qiagen) and ligated into the pCR2.1 cloning vector (Invitrogen). Recombinant plasmids were obtained by standard methodology, including transformation into One Shot *E. coli* cells (Invitrogen), growth of transformed bacterial cells on LB-agar plates containing 100 µg/mL ampicillin (Sigma), growth of bacterial colonies in LB Medium (Life Technologies), and preparation of plasmid DNA using the QIAprep Spin Miniprep Kit (Qiagen). DNA sequence was obtained using cycle sequencing dideoxy terminator reagents (ABI) and synthetic oligonucleotides (see above) reacted in a Gene Amp 9600 thermal cycler (Perkin Elmer) under manufacturer's instructions. Reactions were run and data were generated on an ABI automated sequencer. Sequence data were analyzed using Lasergene software (DNA Star).

**Example 4: Cloning of MORT1 isoforms from brain tissue**

The MORT1 cDNA from human fetal brain was generated using standard PCR techniques. Thermal cycling was performed, using the regime, 1 min at 95C, then 30 cycles (40 sec at 95C, 1 min at 60C, 1 min at 72C), 10 min at 72C, and using the oligonucleotides "5' MORT1" and "3' MORT1" (see above). A human fetal brain (matchmaker) cDNA library (Clontech) and a human adult whole brain (matchmaker) cDNA library (Clontech) were used as DNA templates for PCR. The PCR products of approximately 640 bp were obtained, ligated to the vector pCR II (Invitrogen), and recombinant plasmids were obtained by standard methodology, including transformation into competent One shot *E. coli* cells, growth of transformed bacterial cells on LB-agar plates containing 100 µg/mL ampicillin (Sigma), growth of bacterial colonies in LB Medium (Life Technologies), and preparation of plasmid DNA using by Wizard Plus Minipreps (Promega) or by Qiagen Midiprep (Qiagen). Recombinant plasmids were confirmed by restriction enzyme digestion. The cDNA sequence was determined by dye-deoxy terminator reaction using reagents and protocols from ABI-Perkin-Elmer and the ABI 373A automated sequencer.

The sequence of each of five clones isolated from human fetal brain cDNA matched the published MORT1 sequence. Five clones isolated from human adult brain cDNA were also sequenced. Three of five clones encoded the isoform that contained the deletion of bp 172-192 of the MORT1 coding sequence; the other two clones encoded a single base pair substitution (G173A) relative to the Genbank (X84709) MORT1 sequence.

**Example 5: Cloning of MACH $\alpha$ 1 from NTERA2 cell cDNA**

NTERA2 cells (Stratagene) were grown under conditions to promote terminal differentiation to the neuronal phenotype, including long-term incubation in retinoic acid (Andrews, 1984). Eighteen hours prior to RNA extraction, staurosporine (Calbiochem) was added to the cell culture medium to a final concentration of 100 nM to induce apoptosis. Total RNA extraction was by the BRL RNeasy Total RNA Kit (Life Technologies, Inc.) using reagents and instructions provided by the manufacturer. Reverse transcription of total RNA was performed with the reagents and instructions provided in the BRL Preamplification Kit (Life Technologies, Inc.). Specifically, reverse transcription of 4  $\mu$ g total RNA was primed using oligo(dT).

Oligonucleotides for use in PCR were prepared (on an ABI oligosynthesizer), designed according to the published MACH sequence (Boldin, *et al.*, 1996). A 24 base molecule (MACH 5'  $\alpha$ , 5'-TTT-AAA-AAG-ATG-GAC-TTC-AGC-AGA-3', spanning the translational start codon of MACH  $\alpha$ 1 and other MACH isoforms) was synthesized. Another 24 base molecule (MACH 3'  $\alpha$ , 5'-ATA-GCA-CCA-TCA-ATC-AGA-AGG-GAA-3', complementary to coding sequence, and spanning the stop codon of MACH  $\alpha$ 1 and MACH  $\alpha$ 2. PCR amplification of NTERA2 cDNA was performed using DNA Taq polymerase and other reagents provided by the manufacturer (Life Technologies) and equimolar amounts of oligonucleotides MACH 5'  $\alpha$  and MACH 3'  $\alpha$  (see above). Thermal cycling of the PCRs was in the Gene Amp 2400 thermal cycler (Perkin



Elmer) under the regime, 1 min at 95C, 35 cycles (15 sec at 95C, 30 sec at 50C, 1 min at 72C), 7 min at 72C. Two PCR products, both between 1.4 and 1.5 kb resulted. The larger product (the cDNA corresponding to the MACH  $\alpha$ 1) was agarose gel purified using Qiaex II reagents and instructions provided by the manufacturer (Qiagen) and ligated to the pCR2.1 cloning vector (Invitrogen). Recombinant plasmids were obtained by standard methodology, including transformation into One Shot *E. coli* cells (Invitrogen), growth of transformed bacterial cells on LB-agar plates containing 100  $\mu$ g/mL ampicillin (Sigma), growth of bacterial colonies in LB Medium (Life Technologies,), and preparation of plasmid DNA using the QIAprep Spin Miniprep Kit (Qiagen). DNA sequence was obtained using cycle sequencing dideoxy terminator reagents (ABI) and synthetic oligonucleotides reacted in a Gene Amp 9600 thermal cycler (Perkin Elmer) under manufacturer's instructions. Reactions were run and data were generated on an ABI automated sequencer. Sequence data were analyzed using Lasergene software (DNA Star).

**Example 6: Interaction of MORT1**

**transcript isoforms with MACHa1 C360S**

To investigate the interaction of MORT1 transcript isoforms, a two-hybrid interaction paradigm was first established with MORT1 and MACHa1 C360S. Genes encoding the fusion proteins were generated by cloning MORT1 and MACHa1 C360S cDNA sequences into plasmids containing the coding regions for the domains of the yeast transcriptional activator proteins, Gal4. DNA binding domain (Gal4) fusion are constructed in pAS1 having a 2m origin of replication which is described in Wade-Harper *et al.*, (1993). Gene activation domain (Gal4) fusions are constructed in pACTII (described in Durfee *et al.*, 1993). The yeast system used in this study is as described by Young and Ozenberger (1995).

The MORT1 cDNA is obtained as a NcoI - Bam HI fragment from MORT1-pCRII plasmid and directionally cloned into pAS1 via NcoI and BamHI restrictions sites to generate MORT1-pAS1 plasmid. The MACHa1

wildtype protein is lethal in yeast. Mutation of the cysteine at amino acid position 360 to a serine prevents the lethal phenotype and does not interfere with the protein's ability to interact with MORT1 (Boldin, *et al.*, 1996). The amino acid substitution to generate MACHa1 C360S was obtained using standard PCR techniques (Finney, 1993). Thermal cycling of these reactions was performed under the regime, 1 min at 95C, then 30 cycles (40 sec at 95C, 1 min at 60C, 1 min at 72C), and 10 min at 72C, and using specific oligonucleotides. All oligonucleotides were prepared on an ABI oligosynthesizer and are designed according to published cDNA for human MACH (Boldin *et al.*, 1996; Genbank X98172-X98178). Four oligonucleotides were prepared. A 36 base 5' mutagenic oligonucleotide (oCB1) contained the C360S substitution (TGT to TCT) and a silent mutations at Ala 359 (GCT to GCA) to encrypt a diagnostic restriction enzyme site for SfiI [5'-GTG TTT TTT ATT CAG GCA TCT CAG GGG GAT AAC TAC - 3']. A 36 base 3' mutagenic oligonucleotide (oCB2) [5'-GTA GTT ATC CCC CTG AGA TGC CTG AAT AAA AAA CAC-3'] contained sequence complimentary to oCB1. A 34 base oligonucleotide (oCB3) containing a BamHI site [5'-CGG GAT CCG TAT GGA CTT CAG CAG AAA TCT TTA T-3']. A 38 base 3' oligonucleotide (oCB4) contained a BamHI and Sall restriction enzyme sites [5'-CGG GAT CCG ACG TCG ACT CAA TCA GAA GGG AAG ACA AG-3']. The MACHa1-pCR2.1 plasmid (see example 5, above) was used as template with oligonucleotide pair oCB2 and oCB3 to generate a 1100 basepair fragment, and as template for oligonucleotide pair oCB1 and oCB4 to generate a 360 bp fragment. A subsequent PCR reaction was performed using the 1100 and 360 basepair fragments with oligonucleotide pair oCB3 and oCB4 to generate a 1460 bp fragment which was ligated to pCRII vector (Invitrogen) to generate MACHa1 C360S -pCRII. Recombinant plasmids were obtained by standard methodology, including transformation into One Shot *E. coli* cells (Invitrogen), growth of transformed bacterial cells on LB-agar (Gibco, Inc.)

plates containing either 100 mg/mL ampicillin (Sigma) or 50 mg/mL Kanamycin (Sigma) as described in the manufacturer's (Invitrogen) standard protocol. Growth of bacterial colonies was in LB medium and preparation of plasmid DNA was performed using Qiagen Midiprep (Qiagen) or Wizard Plus Minipreps (Promega). Recombinant plasmids were confirmed restriction enzyme digestion. The cDNA sequence was obtained by dye-deoxy terminator reactions using reagents and protocols from ABI-Perkin-Elmer and the ABI 373A automated sequencer. The cDNA encoding MACHa1 C360S was obtained as a BamHI - Sall fragment and directionally cloned into pACTII via BamHI - XhoI sites to generate MACHa1 C360S - pACTII. DNA of final recombinant vectors were transformed into yeast strain(s) by the lithium acetate method (Rose *et al.*, 1990).

A yeast host strain (CY770) as described in Young and Ozenberger (1995) was transformed with both the MORT1 and MACHa1 C360S fusion constructs (YCB5) or a single fusion construct plus the opposing vector containing unrelated heterologous DNA (YCB5.1 or YCB2.1, respectively). All strains were found to exhibit equal growth on nonselective medium. Strains were then tested for growth on selective medium (*i.e.*, growth medium lacking histidine). Only cells expressing both the MORT1 and MACHa1 C360S fusions are able to grow on selective medium while the strains containing either the MORT1 or the MACHa1 C360S fusion with an unrelated fusion cannot grow (data not shown).

The MORT1 G173A SEQ. (ID NO. 5) was obtained as a NcoI-BamHI fragment from MORT1 G173A-pCRII and directionally cloned into pAS1 to generate MORT1 G173A-pAS1 plasmid. MORT1  $\Delta$ 21 was obtained as a NcoI-BamHI fragment from MORT1 $\Delta$ 21-pCRII and directionally cloned into pAS1 to generated MORT1 $\Delta$ 21-pAS1 plasmid. These plasmids and the MACHa1 C360S -pACTII plasmid were used to transform the yeast host strain, CY770. Strains were generated that

express MORT1 G173A and MACHa1 C360S fusion constructs (YCB16) or the MORT1 G173A fusion construct plus the opposing vector containing unrelated heterologous DNA (YCB16.1). Strains were generated that express MORT1 $\Delta$ 21 and MACHa1 C360S fusion  
5 constructs (YCB9) or the MORT1 $\Delta$ 21 fusion construct plus the opposing vector containing unrelated heterologous DNA (YCB9.1).

Two independent samples of each strain were streaked on standard synthetic medium containing 0, 20 or 40 mM 3-aminotriazole (Figure 2). Plates were incubated at 30C for 3 days. All strains were  
10 found to exhibit equal growth on non-selective medium. Strains were then tested for growth on selective medium (*i.e.*, growth medium lacking histidine). Cells expressing both the MORT1 and MACHa1 C360S fusions, and cells expressing the MORT1 G173A isoform and MACHa1 C360S fusions, were able to grow on selective medium while the strains  
15 containing the MORT1 $\Delta$ 21 and MACHa1 C360S fusions were not able to grow (Figure 2). Negative control strains expressing any of the MORT1 isoform fusions or the MACHa1 C360S fusion with an unrelated fusion cannot grow.

These data suggest that the MORT1 $\Delta$ 21 isoform fusion protein is  
20 impaired in its ability to functionally interact with the MACHa1 C360S fusion protein.

**Example 7: Screen for compounds affecting the interaction  
of MORT1 transcript isoforms with other interacting  
proteins (MACH family members)**

25 The screening methods as described in Young and Ozenberger, WO 95/34646 published December 21, 1995, the whole of which is incorporated herein, are used to identify compounds that affect the interaction of MORT1 transcript isoforms with proteins such as MACHa1 C360S.

30 Low copy number plasmids expressing MORT1 transcript isoforms (MORT1 G173A or MORT1 $\Delta$ 21) and MACHa1 C360S as GAL4 fusion

proteins are constructed to reduce expression of these proteins. cDNA inserts encoding the MORT1 transcript isoforms are subcloned directly from the recombinant pAS1 vector(s) to pUN30AS via NcoI-BamHI fragment to generate either MORT1 G173A-pUN30AS or MORT1 D 21-pUN30AS. The MACHa1 C360S is subcloned directly from the MACHa1 C360S - pCR II as a BamHI-SalI fragment into pUN100ACT via BamHI - XhoI sties to generate MACHa1 C360S - pUN100ACT. These plasmids are transformed into yeast strain CY770 (Young and Ozenberger, 1995) with reporter plasmid pOZ146 (Young and Ozenberger, 1995) to generate yeast strain YCB6. The presence of all three plasmids is necessary to confer the necessary phenotype to enable a rescue screen method as described in Young and Ozenberger (1995).

The yeast strain (YCB18) containing the MORT1 G173A and MACHa1 C360S fusion plasmids plus the reporter plasmid, or the yeast strain (YCB17) containing the MORTD21 and MACHa1 C360S plus the reporter plasmid, forms the basis of a simple primary screen for compounds that disrupt the interaction of the MORT1 transcript isoform gene products and MACHa1 C360S.

**Example 8: Screen for compounds affecting the interaction of MORT1 transcript isoforms and protein containing a death domain motif (TRADD, Fas/APO1 receptor, TNFR)**

The screening methods as described in Young and Ozenberger, WO 95/34646 published December 21, 1995, the whole of which is incorporated herein, are used to identify compounds that affect the interaction of MORT1 transcript isoforms with proteins containing a death domain motif.

Low copy number plasmids expressing MORT1 transcript isoforms (MORT1 G173A, or MORT1Δ21) and the cytoplasmic domain of the TNF receptor containing a death domain motif as GAL4 fusion proteins are constructed to reduce expression of these proteins. cDNA inserts encoding the MORT1 transcript isoforms are subcloned directly from the

MORT1 G173A- pAS1 or the MORT1 $\Delta$ 21-pAS1 recombinant to pUN30AS via NcoI-BamHI fragment to generate either MORT1 G173A-pUN30AS or MORT1 $\Delta$ 21-pUN30AS. The cytoplasmic domain of the TNFR is subcloned directly from TNFcyto-pCR II as an EcoRI fragment into

5 pUN100ACT via EcoRI sites to generate TNFRcyto - pUN100ACT. Plasmids are transformed into the yeast strain CY770 (Young and Ozenberger, 1995) with the reporter plasmid pOZ146 (Young and Ozenberger, 1995). The presence of three plasmids is necessary to confer the necessary phenotype to enable a rescue screen method as

10 described in Young and Ozenberger (1995). The yeast strain (YCB21) containing the MORT1 G173A and TNFRcyto fusion plasmids plus the reporter plasmid, or the yeast strain (YCB20) containing the MORT1 $\Delta$ 21 and TNFRcyto plus the reporter plasmid, forms the basis of a simple primary screen for compounds that disrupt the interaction of the MORT1

15 transcript isoform gene products and a death domain containing protein, TNFR cytoplasmic domain.

## REFERENCES

- Andrews PW.** (1984) Retinoic acid induces neuronal differentiation of a cloned human embryonal carcinoma cell line. *In Vitro Dev Biol* **103**: 285-293
- 5 **Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith J and Struhl K (eds.).** (1993) *Current Protocols in Molecular Biology*, Vol 2. John Wiley & Sons, New York
- Berk AJ, Sharp PA.** (1977) Sizing and mapping of early adenovirus mRNAs by gel electrophoresis of S1 endonuclease-digested hybrids. *Cell*
- 10 **12**: 721-732
- Boldin MP, Varfolomeev EE, Pancer Z, Mett IL, Camonis JH, Wallach D.** (1995) A novel protein that interacts with the death domain of Fas/APO1 contains a sequence motif related to the death domain. *J Biol Chem* **270**: 7795-7798
- 15 **Boldin MP, Goncharov TM, Goltsev YV, Wallace D.** (1996) Involvement of MACH, a novel MORT1/FADD -interacting protease, in Fas/Apo1 - and TNF receptor induced cell death. *Cell* **85**: 803-815
- Chinnaiyan AM, O'Rourke K, Tewari M, Dixit VM.** (1995) FADD, a novel death domain-containing protein, interacts with the death domain of Fas
- 20 and initiates apoptosis. *Cell* **81**: 505-512
- Cordell HJ, Todd JA, Bennett ST, Kawaguchi Y, Farrall M.** (1995) Two-locus maximum lod score analysis of a multifactorial trait: joint consideration of IDDM2 and IDDM4 with IDDM1 in type 1 diabetes. *Am J Hum Genet* **57**: 920-934
- 25 **Coico R (ed.).** (1994) *Current Protocols in Immunology*. John Wiley & Sons, New York
- Davies JL, Kawaguchi Y, Bennett ST, Copeman JB, Cordell HJ, Pritchard LE, Reed PW, Gough SC, Jenkins SC, Palmer SM, et al.** (1994) A genome-wide search for human type 1 diabetes susceptibility genes.
- 30 *Nature* **371**: 130-136

- Durfee T, Becherer K, Chen P-L, Yeh S-H, Yang Y, Kilburn AE, Lee W-H, Elledge SJ. (1993) The retinoblastoma protein associated with the protein phosphatase type 1 catalytic sub-unit. *Genes and Devel.* 7: 555-569
- 5 Farrow SM. (1997) Basis of gene structure and function. *In*: Rumsby G and Farrow M (eds.), *Molecular Endocrinology*. BIOS Scientific Publishers, Ltd, Oxford, pp 1-22
- Finney M. (1993) The polymerase chain reaction. *In*: Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith J and Struhl K (eds.),
- 10 *Current Protocols in Molecular Biology*, Vol 2. John Wiley & Sons, New York, Chapter 15
- Fisher GH, Rosenberg FJ, Straus SE, Dale JK, Middleton LA, Lin AY, Strober W, Lenardo MJ, Puck JM. (1995) Dominant interfering Fas gene mutations impair apoptosis in a human autoimmune lymphoproliferative
- 15 syndrome. *Cell* 81: 935-946
- Freshney RI. (1987) *Culture of Animal Cells*. A R Liss, New York
- Glover DN (ed.). (1985) *DNA Cloning*, Vol 1 and 2. IRL Press, Oxford
- Griffin HG, Griffin AM (eds.). (1994) *PCR Technology: Current Innovations*. CRC Press, Boca Raton, FL
- 20 Hanes BD, Higgins SJ. (eds.) (1984) *Transcription and Translation*. IRL Press, Oxford
- Hashimoto L, Habita C, Beressi JP, Delepine M, Bessé C, Cambon-Thomsen A, Deschamps I, Rotter JI, Djoulah S, James MR, et al. (1994) Genetic mapping of a susceptibility locus for insulin-dependent diabetes
- 25 mellitus on chromosome 11q. *Nature* 371: 161-164
- Henkart, PA. (1996) ICE family proteases: mediators of all apoptotic cell death? *Immunity* 4: 195-201
- Kim PKM, Dutra AS, Chandrasekharappa SC, Puck JM. (1996) Genomic structure and mapping of human FADD, an intracellular mediator of
- 30 lymphocyte apoptosis. *J Immunol* 157: 5461-5466.



- Kischkel FC, Hellbardt S, Behrmann I, Germer M, Pawlita M, Krammer PH, Peter ME. (1995) Cytotoxicity-dependent APO-1 (Fas/CD95)-associated proteins form a death-inducing signaling complex (DISC) with the receptor. *EMBO J* 14: 5579-5588
- 5 Lafage M, Pedeutour F, Marchetto S, Simonetti J, Prosperi MT, Gaudray P, Birnbaum D. (1992) Fusion and amplification of two originally non-syntenic chromosomal regions in a mammary carcinoma cell line. *Genes Chromosomes Cancer* 5: 40-49
- Lee JJ, Costlow NA. (1987) A molecular titration assay to measure transcript prevalence levels. *Methods Enzymol* 152: 633-648
- 10 Lewin B. (1994) *Genes V*. Oxford University and Cell Press, Oxford.
- Nagata S. (1994) Fas and Fas ligand: a death factor and its receptor. *Adv Immunol* 57: 129-144
- Perbal B. (1988) *A Practical Guide to Molecular Cloning*, 2nd ed. John Wiley & Sons, New York
- 15 Rieux-Laucat F, Le Deist F, Hivroz C, Roberts IA, Debatin KM, Fischer A, de Villartay JP. (1995) Mutations in Fas associated with human lymphoproliferative syndrome and autoimmunity. *Science* 268: 1347-1349
- 20 Rose, MD, Winston, F, Hieter, P. (1990) *Methods in yeast genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sambrook J, Fritsch EF, Maniatis T. (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY
- 25 Schuurin E. (1995) The involvement of the chromosome 11q13 region in human malignancies: cyclin D1 and EMS1 are two new candidate oncogenes--a review. *Gene* 159: 83-96
- Schuuring E, Verhoeven E, van Tinteren H, Peterse JL, Nunnink B, Thunnissen FB, Devilee P, Cornelisse CJ, van de Vijver MJ, Mooi WJ, et al. (1992) Amplification of genes within the chromosome 11q13 region
- 30

is indicative of poor prognosis in patients with operable breast cancer.

Cancer Res 52: 5229-5234

**Szepietowski P, Perucca-Lostanlen D, Grosgeorge J, LePaslier D, Brownstein BH, Carle GF, Gaudray P.** (1995) Description of a 700-kb

5 yeast artificial chromosome contig containing the BCL1 translocation breakpoint region at 11q13. Cytogenet Cell Genet 69: 101-107

**Thompson CB.** (1995) Apoptosis in the pathogenesis and treatment of disease. Science 267: 1456-1462

10 **Tisch R, McDevitt H.** (1996) Insulin-dependent diabetes mellitus. Cell 85: 291-297

**Wade-Harper J, Adami GR, Wei N, Keyomarsk K, Elledge SJ.** (1993) The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. Cell 75: 801-816

15 **Young, KH, Ozenberger, BA.** (1995) Novel cell systems having specific interaction of peptide binding pairs. WO 95/34646. December 21.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Wood, Andrew T  
Bingham, Brendan W  
Young, Kathleen H  
Birsan, Camelia
- (ii) TITLE OF INVENTION: Neuronal MORT1 Isoforms
- (iii) NUMBER OF SEQUENCES: 6
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Andrea C. Walsh
  - (B) STREET: One Campus Drive
  - (C) CITY: Parsippany
  - (D) STATE: New Jersey
  - (E) COUNTRY: USA
  - (F) ZIP: 07054
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: US
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Walsh, Andrea C.
  - (B) REGISTRATION NUMBER: 34,988
  - (C) REFERENCE/DOCKET NUMBER: AHP-97147
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (973) 683-2169
  - (B) TELEFAX: (973) 683-4117

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 606 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 1..606

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG GAC CCG TTC CTG GTG CTG CTG CAC TCG GTG TCG TCC AGC CTG TCG	48
Met Asp Pro Phe Leu Val Leu Leu His Ser Val Ser Ser Ser Leu Ser	
1 5 10 15	
AGC AGC GAG CTG ACC GAG CTC AAG TTC CTA TGC CTC GGG CGC GTG GGC	96
Ser Ser Glu Leu Thr Glu Leu Lys Phe Leu Cys Leu Gly Arg Val Gly	
20 25 30	
AAG CGC AAG CTG GAG CGC GTG CAG AGC GGC CTA GAC CTC TTC TCC ATG	144
Lys Arg Lys Leu Glu Arg Val Gln Ser Gly Leu Asp Leu Phe Ser Met	
35 40 45	
CTG CTG GAG CAG AAC GAC CTG GAG CCC GAG CTG CTC GCC TCC CTG CGG	192
Leu Leu Glu Gln Asn Asp Leu Glu Pro Glu Leu Leu Ala Ser Leu Arg	
50 55 60	
CGC CAC GAC CTG CTG CGG CGC GTC GAC GAC TTC GAG GCG GGG GCG GCG	240
Arg His Asp Leu Leu Arg Arg Val Asp Asp Phe Glu Ala Gly Ala Ala	
65 70 75 80	
GCC GGG GCC GCG CCT GGG GAA GAA GAC CTG TGT GCA GCA TTT AAC GTC	288
Ala Gly Ala Ala Pro Gly Glu Glu Asp Leu Cys Ala Ala Phe Asn Val	
85 90 95	
ATA TGT GAT AAT GTG GGG AAA GAT TGG AGA AGG CTG GCT CGT CAG CTC	336
Ile Cys Asp Asn Val Gly Lys Asp Trp Arg Arg Leu Ala Arg Gln Leu	
100 105 110	
AAA GTC TCA GAC ACC AAG ATC GAC AGC ATC GAG GAC AGA TAC CCC CGC	384
Lys Val Ser Asp Thr Lys Ile Asp Ser Ile Glu Asp Arg Tyr Pro Arg	
115 120 125	
AAC CTG ACA GAG CGT GTG CGG GAG TCA CTG AGA ATC TGG AAG AAC ACA	432

27

Asn	Leu	Thr	Glu	Arg	Val	Arg	Glu	Ser	Leu	Arg	Ile	Trp	Lys	Asn	Thr		
130						135					140						
GAG	AAG	GAG	AAC	GCA	ACA	GTG	GCC	CAC	CTG	GTG	GGG	GCT	CTC	AGG	TCC		480
Glu	Lys	Glu	Asn	Ala	Thr	Val	Ala	His	Leu	Val	Gly	Ala	Leu	Arg	Ser		
145					150					155				160			
TGC	CAG	ATG	AAC	CTG	GTG	GCT	GAC	CTG	GTA	CAA	GGG	GTT	CAG	CAG	GCC		528
Cys	Gln	Met	Asn	Leu	Val	Ala	Asp	Leu	Val	Gln	Gly	Val	Gln	Gln	Ala		
				165					170					175			
CGT	GAC	CTC	CAG	AAC	AGG	AGT	GGG	GCC	ATG	TCC	CCG	ATG	TCA	TGG	AAC		576
Arg	Asp	Leu	Gln	Asn	Arg	Ser	Gly	Ala	Met	Ser	Pro	Met	Ser	Trp	Asn		
			180				185						190				
TCA	GAC	GCA	TCT	ACC	TCC	GAA	GCG	TCC	TGA								606
Ser	Asp	Ala	Ser	Thr	Ser	Glu	Ala	Ser	*								
			195				200										

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 202 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Asp	Pro	Phe	Leu	Val	Leu	Leu	His	Ser	Val	Ser	Ser	Ser	Leu	Ser		
1				5					10					15			
Ser	Ser	Glu	Leu	Thr	Glu	Leu	Lys	Phe	Leu	Cys	Leu	Gly	Arg	Val	Gly		
		20					25						30				
Lys	Arg	Lys	Leu	Glu	Arg	Val	Gln	Ser	Gly	Leu	Asp	Leu	Phe	Ser	Met		
		35					40					45					
Leu	Leu	Glu	Gln	Asn	Asp	Leu	Glu	Pro	Glu	Leu	Leu	Ala	Ser	Leu	Arg		
		50				55						60					
Arg	His	Asp	Leu	Leu	Arg	Arg	Val	Asp	Asp	Phe	Glu	Ala	Gly	Ala	Ala		
	65				70					75					80		
Ala	Gly	Ala	Ala	Pro	Gly	Glu	Glu	Asp	Leu	Cys	Ala	Ala	Phe	Asn	Val		
			85					90						95			
Ile	Cys	Asp	Asn	Val	Gly	Lys	Asp	Trp	Arg	Arg	Leu	Ala	Arg	Gln	Leu		
		100					105						110				
Lys	Val	Ser	Asp	Thr	Lys	Ile	Asp	Ser	Ile	Glu	Asp	Arg	Tyr	Pro	Arg		
		115					120					125					
Asn	Leu	Thr	Glu	Arg	Val	Arg	Glu	Ser	Leu	Arg	Ile	Trp	Lys	Asn	Thr		
		130				135						140					

28

Glu Lys Glu Asn Ala Thr Val Ala His Leu Val Gly Ala Leu Arg Ser  
 145 150 155 160

Cys Gln Met Asn Leu Val Ala Asp Leu Val Gln Gly Val Gln Gln Ala  
 165 170 175

Arg Asp Leu Gln Asn Arg Ser Gly Ala Met Ser Pro Met Ser Trp Asn  
 180 185 190

Ser Asp Ala Ser Thr Ser Glu Ala Ser \*  
 195 200

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 606 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..606

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATG GAC CCG TTC CTG GTG CTG CTG CAC TCG GTG TCG TCC AGC CTG TCG	48
Met Asp Pro Phe Leu Val Leu Leu His Ser Val Ser Ser Ser Leu Ser	
1 5 10 15	
AGC AGC GAG CTG ACC GAG CTC AAG TTC CTA TGC CTC GGG CGC GTG GGC	96
Ser Ser Glu Leu Thr Glu Leu Lys Phe Leu Cys Leu Gly Arg Val Gly	
20 25 30	
AAG CGC AAG CTG GAG CGC GTG CAG AGC GGC CTA GAC CTC TTC TCC ATG	144
Lys Arg Lys Leu Glu Arg Val Gln Ser Gly Leu Asp Leu Phe Ser Met	
35 40 45	
CTG CTG GAG CAG AAC GAC CTG GAG CCC GAG CTG CTC GCC TCC CTG CGG	192
Leu Leu Glu Gln Asn Asp Leu Glu Pro Glu Leu Leu Ala Ser Leu Arg	
50 55 60	
CGC CAC GAC CTG CTG CGG CGC GTC GAC GAC TTC GAG GCG GGG GCG GCA	240
Arg His Asp Leu Leu Arg Arg Val Asp Asp Phe Glu Ala Gly Ala Ala	
65 70 75 80	
GCC GGG GCC GCG CCT GGG GAA GAA GAC CTG TGT GCA GCA TTT AAC GTC	288
Ala Gly Ala Ala Pro Gly Glu Glu Asp Leu Cys Ala Ala Phe Asn Val	
85 90 95	
ATA TGT GAT AAT GTG GGG AAA GAT TGG AGA AGG CTG GCT CGT CAG CTC	336
Ile Cys Asp Asn Val Gly Lys Asp Trp Arg Arg Leu Ala Arg Gln Leu	
100 105 110	

29

AAA GTC TCA GAC ACC AAG ATC GAC AGC ATC GAG GAC AGA TAC CCC CGC	384
Lys Val Ser Asp Thr Lys Ile Asp Ser Ile Glu Asp Arg Tyr Pro Arg	
115 120 125	
AAC CTG ACA GAG CGT GTG CGG GAG TCA CTG AGA ATC TGG AAG AAC ACA	432
Asn Leu Thr Glu Arg Val Arg Glu Ser Leu Arg Ile Trp Lys Asn Thr	
130 135 140	
GAG AAG GAG AAC GCA ACA GTG GCC CAC CTG GTG GGG GCT CTC AGG TCC	480
Glu Lys Glu Asn Ala Thr Val Ala His Leu Val Gly Ala Leu Arg Ser	
145 150 155 160	
TGC CAG ATG AAC CTG GCG GCT GAC CTG GTA CAA GAG GTT CAG CAG GCC	528
Cys Gln Met Asn Leu Ala Ala Asp Leu Val Gln Glu Val Gln Gln Ala	
165 170 175	
CGT GAC CTC CAG AAC AGG AGT GGG GCC ATG TCC CCG ATG TCA TGG AAC	576
Arg Asp Leu Gln Asn Arg Ser Gly Ala Met Ser Pro Met Ser Trp Asn	
180 185 190	
TCA GAC GCA TCT ACC TCC GAA GCG TCC TGA	606
Ser Asp Ala Ser Thr Ser Glu Ala Ser *	
195 200	

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 202 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Asp Pro Phe Leu Val Leu Leu His Ser Val Ser Ser Ser Leu Ser	
1 5 10 15	
Ser Ser Glu Leu Thr Glu Leu Lys Phe Leu Cys Leu Gly Arg Val Gly	
20 25 30	
Lys Arg Lys Leu Glu Arg Val Gln Ser Gly Leu Asp Leu Phe Ser Met	
35 40 45	
Leu Leu Glu Gln Asn Asp Leu Glu Pro Glu Leu Leu Ala Ser Leu Arg	
50 55 60	
Arg His Asp Leu Leu Arg Arg Val Asp Asp Phe Glu Ala Gly Ala Ala	
65 70 75 80	
Ala Gly Ala Ala Pro Gly Glu Glu Asp Leu Cys Ala Ala Phe Asn Val	
85 90 95	
Ile Cys Asp Asn Val Gly Lys Asp Trp Arg Arg Leu Ala Arg Gln Leu	
100 105 110	
Lys Val Ser Asp Thr Lys Ile Asp Ser Ile Glu Asp Arg Tyr Pro Arg	

30

115		120		125
Asn Leu Thr Glu Arg Val Arg Glu Ser Leu Arg Ile Trp Lys Asn Thr				
130		135		140
Glu Lys Glu Asn Ala Thr Val Ala His Leu Val Gly Ala Leu Arg Ser				
145		150		155
				160
Cys Gln Met Asn Leu Ala Ala Asp Leu Val Gln Glu Val Gln Gln Ala				
		165		170
				175
Arg Asp Leu Gln Asn Arg Ser Gly Ala Met Ser Pro Met Ser Trp Asn				
		180		185
				190
Ser Asp Ala Ser Thr Ser Glu Ala Ser *				
195		200		

## (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 627 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..627

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG GAC CCG TTC CTG GTG CTG CTG CAC TCG GTG TCG TCC AGC CTG TCG	48
Met Asp Pro Phe Leu Val Leu Leu His Ser Val Ser Ser Ser Leu Ser	
1 5 10 15	
AGC AGC GAG CTG ACC GAG CTC AAG TTC CTA TGC CTC GGG CGC GTG GGC	96
Ser Ser Glu Leu Thr Glu Leu Lys Phe Leu Cys Leu Gly Arg Val Gly	
20 25 30	
AAG CGC AAG CTG GAG CGC GTG CAG AGC GGC CTA GAC CTC TTC TCC ATG	144
Lys Arg Lys Leu Glu Arg Val Gln Ser Gly Leu Asp Leu Phe Ser Met	
35 40 45	
CTG CTG GAG CAG AAC GAC CTG GAG CCC GAG CAC ACC GAG CTC CTG CGC	192
Leu Leu Glu Gln Asn Asp Leu Glu Pro Glu His Thr Glu Leu Leu Arg	
50 55 60	
GAG CTG CTC GCC TCC CTG CGG CGC CAC GAC CTG CTG CGG CGC GTC GAC	240
Glu Leu Leu Ala Ser Leu Arg Arg His Asp Leu Leu Arg Arg Val Asp	
65 70 75 80	
GAC TTC GAG GCG GGG GCG GCA GCC GGG GCC GCG CCT GGG GAA GAA GAC	288
Asp Phe Glu Ala Gly Ala Ala Ala Gly Ala Ala Pro Gly Glu Glu Asp	
85 90 95	



CTG TGT GCA GCA TTT AAC GTC ATA TGT GAT AAT GTG GGG AAA GAT TGG Leu Cys Ala Ala Phe Asn Val Ile Cys Asp Asn Val Gly Lys Asp Trp 100 105 110	336
AGA AGG CTG GCT CGT CAG CTC AAA GTC TCA GAC ACC AAG ATC GAC AGC Arg Arg Leu Ala Arg Gln Leu Lys Val Ser Asp Thr Lys Ile Asp Ser 115 120 125	384
ATC GAG GAC AGA TAC CCC CGC AAC CTG ACA GAG CGT GTG CGG GAG TCA Ile Glu Asp Arg Tyr Pro Arg Asn Leu Thr Glu Arg Val Arg Glu Ser 130 135 140	432
CTG AGA ATC TGG AAG AAC ACA GAG AAG GAG AAC GCA ACA GTG GCC CAC Leu Arg Ile Trp Lys Asn Thr Glu Lys Glu Asn Ala Thr Val Ala His 145 150 155 160	480
CTG GTG GGG GCT CTC AGG TCC TGC CAG ATG AAC CTG GCG GCT GAC CTG Leu Val Gly Ala Leu Arg Ser Cys Gln Met Asn Leu Ala Ala Asp Leu 165 170 175	528
GTA CAA GAG GTT CAG CAG GCC CGT GAC CTC CAG AAC AGG AGT GGG GCC Val Gln Glu Val Gln Gln Ala Arg Asp Leu Gln Asn Arg Ser Gly Ala 180 185 190	576
ATG TCC CCG ATG TCA TGG AAC TCA GAC GCA TCT ACC TCC GAA GCG TCC Met Ser Pro Met Ser Trp Asn Ser Asp Ala Ser Thr Ser Glu Ala Ser 195 200 205	624
TGA *	627

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 209 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Asp Pro Phe Leu Val Leu Leu His Ser Val Ser Ser Ser Leu Ser 1 5 10 15
Ser Ser Glu Leu Thr Glu Leu Lys Phe Leu Cys Leu Gly Arg Val Gly 20 25 30
Lys Arg Lys Leu Glu Arg Val Gln Ser Gly Leu Asp Leu Phe Ser Met 35 40 45
Leu Leu Glu Gln Asn Asp Leu Glu Pro Glu His Thr Glu Leu Leu Arg 50 55 60
Glu Leu Leu Ala Ser Leu Arg Arg His Asp Leu Leu Arg Arg Val Asp

65 70 32 75 80  
Asp Phe Glu Ala Gly Ala Ala Ala Gly Ala Ala Pro Gly Glu Glu Asp  
85 90 95  
Leu Cys Ala Ala Phe Asn Val Ile Cys Asp Asn Val Gly Lys Asp Trp  
100 105 110  
Arg Arg Leu Ala Arg Gln Leu Lys Val Ser Asp Thr Lys Ile Asp Ser  
115 120 125  
Ile Glu Asp Arg Tyr Pro Arg Asn Leu Thr Glu Arg Val Arg Glu Ser  
130 135 140  
Leu Arg Ile Trp Lys Asn Thr Glu Lys Glu Asn Ala Thr Val Ala His  
145 150 155 160  
Leu Val Gly Ala Leu Arg Ser Cys Gln Met Asn Leu Ala Ala Asp Leu  
165 170 175  
Val Gln Glu Val Gln Gln Ala Arg Asp Leu Gln Asn Arg Ser Gly Ala  
180 185 190  
Met Ser Pro Met Ser Trp Asn Ser Asp Ala Ser Thr Ser Glu Ala Ser  
195 200 205

\*

**What is Claimed is:**

1. A composition comprising an isolated polynucleotide from NTERA2 cells selected from the group consisting of:


- (a) a polynucleotide comprising the MORT1Δ21 nucleotide sequence set forth in SEQ. ID NO. 1, deposited under accession number ATCC 209013;
- (b) a polynucleotide comprising the nucleotide sequence encoding a protein that interacts with the death domain of Fas/APO1 or fragments thereof;
- (c) a polynucleotide encoding a protein that interacts with MACH  $\alpha$ 1 or other members of the ICE/Ced3 (Caspase) family of proteins or fragments thereof;
- (d) a polynucleotide which is an allelic variant of the polynucleotide of (a)-(c) above; and
- (e) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(d).

2. A composition comprising an isolated polynucleotide from human brain selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence set forth as in SEQ. ID NO. 3, deposited under accession number ATCC 209018;
- (b) a polynucleotide encoding a protein that interacts with the death domain of Fas/APO1 or fragments thereof;
- (c) a polynucleotide encoding a protein that interacts with MACH  $\alpha$ 1 or other members of the ICE/Ced3 (Caspase) family of proteins or fragments thereof;
- (d) a polynucleotide which is an allelic variant of the polynucleotide of (a)-(c) above; and

- (e) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(d).
3. A composition comprising an isolated polynucleotide from human brain selected from the group consisting of:
- (a) a polynucleotide comprising the nucleotide sequence set forth in SEQ. ID NO. 5, deposited under accession number ATCC 209019;
  - (b) a polynucleotide encoding a protein that interacts with the death domain of Fas/APO1 or fragments thereof;
  - (c) a polynucleotide encoding a protein that interacts with MACH  $\alpha$ 1 or other members of the ICE/Ced3 (Caspase) family of proteins or fragments thereof;
  - (d) a polynucleotide which is an allelic variant of the polynucleotide of (a)-(c) above; and
  - (e) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(d).
4. An isolated MORT1 nucleic acid isoform, wherein the isoform is a human NTERA2 neuronal cell or human brain isoform encoding a protein comprising the amino acid sequences selected from the group consisting of SEQ. ID NO. 2, SEQ. ID NO. 4 AND SEQ. ID NO. 5.
5. An isolated MORT1 nucleic acid isoform comprising the amino acid sequence of SEQ. ID NO. 2.
6. An isolated MORT1 nucleic acid isoform comprising the amino acid sequence of SEQ. ID NO. 4.
7. An isolated MORT1 nucleic acid isoform comprising the amino acid sequence of SEQ. ID NO. 6.

1/2



SC -his, ura, leu, trp

FIG.1

2/2




FIG.2

## INTERNATIONAL SEARCH REPORT

Interr national application No

PCT/US 98/07439

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 C12N15/12 C07K14/47 C07K14/715

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
 IPC 6 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 31603 A (UNIV MICHIGAN ;DIXIT VISHVA M (US); ROURKE KAREN O (US)) 10 October 1996 see page 21, paragraph 3; claims 1-14; figure 2; example 1	2
Y	WO 96 18641 A (YEDA RES & DEV ;WEINWURZEL HENRY (IL); WALLACH DAVID (IL); BOLDIN) 20 June 1996 see abstract; claims 1-28; example 1; table 1	1-4

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 September 1998

Date of mailing of the international search report

25/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Gurdjian, D

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/07439

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>WANKER, ERICH E. ET AL: "HIP-I: a huntingtin interacting protein isolated by the yeast two-hybrid system"  HUM. MOL. GENET. (OCTOBER 25-30 1997),  6(3), 487-495 CODEN: HMGE55;ISSN:  0964-6906, XP002077553  see abstract  see page 493, left-hand column, paragraph 2 - page 494, left-hand column, paragraph 2</p>	1-4
X	<p>CHINNAIYAN A M ET AL: "FADD, A NOVEL DEATH DOMAIN-CONTAINING PROTEIN, INTERACTS WITH THE DEATH DOMAIN OF FAS AND INITIATES APOPTOSIS"  CELL,  vol. 81, no. 4, 19 May 1995, pages  505-512, XP002015439  see the whole document</p>	7
X	<p>WO 96 25941 A (PANCER ZEEV ;GONCHAROV TANYA M (IL); METT IGOR (IL); WALLACH DAVID) 29 August 1996  see claims 1-36; figure SEQ.ID.5</p>	5,6
A	<p>BREDESEN DE: "KEEPING NEURONS ALIVE - THE MOLECULAR CONTROL OF APOPTOSIS.1."  NEUROSCIENTIST, 1996, 2, 181-190,  XP002077555  see abstract; table 4</p>	1-7
P, X	<p>BINGHAM B ET AL: "Human neuronal sequence variants of the mediator of apoptosis, MORT1"  SOCIETY FOR NEUROSCIENCE ABSTRACTS, 23 (1-2). 896., October 1997, XP002077554  see the whole document</p>	1-4



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/07439

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9631603 A	10-10-1996	AU 5301296 A	23-10-1996
WO 9618641 A	20-06-1996	AU 4602296 A	03-07-1996
		CA 2207815 A	20-06-1996
		FI 972457 A	10-06-1997
		NO 972716 A	11-07-1997
		AU 6180596 A	18-02-1997
		NO 980198 A	09-03-1998
		WO 9703998 A	06-02-1997
WO 9625941 A	29-08-1996	AU 5133296 A	11-09-1996
		CA 2213484 A	29-08-1996
		EP 0813419 A	29-12-1997

Form PCT/ISA/210 (patent family annex) (July 1992)

